

Q	Marking Instructions	AO	Marks	Typical Solution
1	Circles correct answer	1.1b	B1	4
<b>Total</b>			<b>1</b>	

Q	Marking Instructions	AO	Marks	Typical Solution
2	Circles correct answer	2.3	B1	$n = 6$
<b>Total</b>			<b>1</b>	

Q	Marking Instructions	AO	Marks	Typical Solution
3(a)	Substitutes $x = -1$ or $x = 3$ into $f(x)$ to obtain one equation or uses identity eg $f(x) \equiv (x + 1)(x - 3)(ax + b)$	3.1a	M1	$x = -1 \quad -p - 3 + 8 + q = 0$ $x = 3 \quad 27p - 27 - 24 + q = 0$
	Obtains two correct equations by substitution method ACF or obtains $a = 2, b = 1$	1.1b	A1	$p = 2$ and $q = -3$
	Solves to find $p$ and $q$ CAO	1.1b	A1	
3(b)	Uses inspection, division by quadratic factor or repeated division or finds third root $x = -\frac{1}{2}$ PI by $(x + \frac{1}{2})$	1.1a	M1	$(x + 1)(x - 3) = x^2 - 2x - 3$ $(x^2 - 2x - 3)(2x + 1)$ $(x + 1)(x - 3)(2x + 1)$
	Completes factorisation	1.1b	A1	
<b>Total</b>			<b>5</b>	

Q	Marking Instructions	AO	Marks	Typical Solution
4	Multiplies by $\frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} + \sqrt{2}}$	AO1.1a	M1	$\frac{\sqrt{6}}{\sqrt{3} - \sqrt{2}} \times \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} + \sqrt{2}}$
	Correctly evaluates denominator to get $3 - 2$ or $1$	AO1.1b	A1	$= \frac{\sqrt{18} + \sqrt{12}}{3 - 2}$
	Evaluates numerator, one term correct $\sqrt{18}$ or $\sqrt{12}$ or $3\sqrt{2}$ or $2\sqrt{3}$	AO1.1b	A1	$\frac{\sqrt{18} + \sqrt{12}}{1}$
	Completes solution CAO	AO2.1	R1	$= \sqrt{9 \times 2} + \sqrt{4 \times 3}$ $= 3\sqrt{2} + 2\sqrt{3}$
<b>Total</b>			<b>4</b>	