




Q1. The diagram shows a eukaryotic cell.



(a) Complete the table by giving the letter labelling the organelle that matches the function.

Function of organelle	Letter
Protein synthesis	
Modifies protein (for example, adds carbohydrate to protein)	
Aerobic respiration	

(3)

(b) Use the scale bar in the diagram above to calculate the magnification of the drawing. Show your working.

Answer = .....

(2)

(Total 5 marks)



Q2.

(a) What is a monomer?

-----  
-----  
-----  
-----  
-----

(1)

(b) Lactulose is a disaccharide formed from one molecule of galactose and one molecule of fructose.

Other than both being disaccharides, give one similarity and one difference between the structures of lactulose and lactose.

Similarity -----

-----  
-----  
-----  
-----

Difference -----

-----  
-----  
-----  
-----

(2)



- (c) Following digestion and absorption of food, the undigested remains are processed to form faeces in the parts of the intestine below the ileum.

The faeces of people with constipation are dry and hard. Constipation can be treated by drinking lactulose. Lactulose is soluble, but is not digested or absorbed in the human intestine.

Use your knowledge of water potential to suggest why lactulose can be used to help people suffering from constipation.

-----  
-----  
-----  
-----  
-----  
-----  
-----  
-----  
-----  
-----

(2)

- (d) Lactulose can also be used to treat people who have too high a concentration of hydrogen ions ( $H^+$ ) in their blood.

The normal range for blood  $H^+$  concentration is  $3.55 \times 10^{-8}$  to  $4.47 \times 10^{-8} \text{ mol dm}^{-3}$

A patient was found to have a blood  $H^+$  concentration of  $2.82 \times 10^{-7} \text{ mol dm}^{-3}$

Calculate the minimum percentage decrease required to bring the patient's blood  $H^+$  concentration into the normal range.

Answer = -----

(2)

(Total 7 marks)



**Q3.**

- (a) Draw **and** label a single DNA nucleotide.

(2)

- (b) Give **two** features of DNA **and** explain how each one is important in the semi-conservative replication of DNA.

1. ....

.....

.....

2. ....

.....

.....

(2)

- (c) Replication of mitochondrial DNA (mtDNA) is different from that of nuclear DNA.

The replication of the second strand of mtDNA **only** starts after two-thirds of the first strand of mtDNA has been copied.

A piece of mtDNA is 16 500 base pairs long and is replicated at a rate of 50 nucleotides per second.

Tick (✓) the box that shows how long it would take to copy this mtDNA.

**A** 330 seconds

**B** 440 seconds

**C** 550 seconds

**D** 660 seconds

(1)

(Total 5 marks)



### Q4.

A scientist used grasshoppers to investigate the effect of composition of air on breathing rate in insects. He changed the composition of air they breathed in by varying the concentrations of oxygen and carbon dioxide.

The scientist collected 20 mature grasshoppers from a meadow. He placed the grasshoppers in a small chamber where he could adjust and control the composition of air surrounding them. The small chamber restricted the movement of the grasshoppers.

His results for three of the grasshoppers are shown in the table below in the form in which he presented them.

		Percentage of oxygen and carbon dioxide in different types of air breathed in by grasshoppers			
		A Air from atmosphere	B Pure oxygen	C Gas mixture 1	D Gas mixture 2
Gas	Oxygen	20.9	100.0	91.0	84.0
	Carbon dioxide	0.1	0.0	9.0	16.0
Breathing rate of grasshopper in different types of air / breaths per minute	Grasshopper 1	53	11	99	107
	Grasshopper 2	48	25	88	99
	Grasshopper 3	61	13	96	93

- (a) The percentages of oxygen and carbon dioxide in Column **A** do **not** add up to 100% but in columns **C** and **D** they do. Suggest **two** reasons for this difference.

1. ....
- .....
2. ....
- .....

(2)



- (b) Use all the data to describe the effect of concentration of carbon dioxide on the breathing rate of grasshoppers.

-----  
-----  
-----  
-----  
-----  
-----

(3)

- (c) One of the different types of air was similar to the air in the meadow where the grasshoppers were collected. It provides data that might be used to calculate a mean breathing rate for grasshoppers in the meadow.

- (i) Use the data to estimate the mean breathing rate of the three grasshoppers in the meadow. Show your working.

Mean breathing rate = ----- breaths per minute

(2)

- (ii) The estimate does not provide a reliable value for the mean breathing rate of all insect species in the meadow. Other than being an estimate, suggest and explain **three** reasons why this value would **not** be reliable.

1. -----  
-----  
-----

2. -----  
-----  
-----

3. -----  
-----  
-----

(3)

(Total 10 marks)



**Q5.**

Farmers use artificial fertilisers to maintain or increase yield from grain-producing crop plants such as wheat.

- (a) Artificial fertiliser is used to replace mineral ions removed from the land when crops are harvested. One of the mineral ions is nitrate.


Give **two** examples of biological molecules containing nitrogen that would be removed when a crop is harvested.

1. ....
2. ....

(2)

- (b) Scientists investigated changes in the use of artificial fertiliser in India between 1970 and 2005. They also investigated changes in the **fertiliser response ratio**. This ratio shows how many kg of grain are produced for each kg of fertiliser used.

The graph shows their results in the form the scientists presented them. (A hectare is a unit of area commonly used in agriculture)



Use these data to calculate the difference in the mass of grain produced per hectare in 1970 compared with 2005.

Show your working.

Difference ..... kg hectare<sup>-1</sup>

(2)



- (c) Use the data in the graph above to evaluate the use of artificial fertilisers on grain-producing crops in India.

-----

-----

-----

-----

-----

-----

**(2)**  
**(Total 6 marks)**

**Q6.**

- (a) Describe the relationship between size and surface area to volume ratio of organisms.

-----

-----

-----

-----

-----

**(1)**

- (b) A scientist calculated the surface area of a large number of frog eggs. He found that the mean surface area was  $9.73 \text{ mm}^2$ . Frog eggs are spherical.

The surface area of a sphere is calculated using this equation

$$\text{Surface area} = 4 \pi r^2 \text{ (where } r \text{ is the radius of a sphere)} \quad \pi = 3.14$$

Use this equation to calculate the mean diameter of a frog egg.

Show your working.

Diameter = ----- mm

**(2)**





The scientist calculated the ratio of surface area to mass for eggs, tadpoles and frogs. He also determined the mean rate of oxygen uptake by tadpoles and frogs.

His results are shown in the table.

Stage of frog development	Ratio of surface area to mass	Mean rate of oxygen uptake / $\mu\text{mol g}^{-1} \text{h}^{-1}$
Egg	2904 : 1	no information
Tadpole	336 : 1	5.7
Adult	166 : 1	1.3

- (c) The scientist used units of  $\mu\text{mol g}^{-1} \text{h}^{-1}$  for the rate of oxygen uptake.

Suggest why he used  $\mu\text{mol}$  in these units.

-----  
 -----  
 -----  
 -----  
 -----

(1)

- (d) The scientist decided to use the ratio of surface area to mass, rather than the ratio of surface area to volume. He made this decision for practical reasons.

Suggest **one** practical advantage of measuring the masses of frog eggs, tadpoles and adults, compared with measuring their volumes.

-----  
 -----  
 -----  
 -----

(1)

- (e) Explain why oxygen uptake is a measure of metabolic rate in organisms.

-----  
 -----  
 -----  
 -----  
 -----



- (f) A student who looked at these results said that they could not make a conclusion about the relationship between stage of development and metabolic rate.

Use information in the table to explain reasons why they were unable to make a conclusion.

-----

-----

-----

-----

-----

-----

-----

-----


(3)  
(Total 9 marks)



**Q7.**

Scientists investigated how the concentration of protein in blood plasma changes in people between the ages of 60 and 95.

The graph shows the scientists' results. The bars show  $\pm 1$  standard deviation.



- (a) What is the difference between males and females in the fall in mean concentration of protein in blood plasma between 60 and 95 years?

Answer = \_\_\_\_\_ g dm<sup>-3</sup> (1)

- (b) Use the graph above to calculate the rate of change of the mean concentration of protein in the blood plasma of males between the ages of 60 and 95.

Show your working.

Answer = \_\_\_\_\_ g dm<sup>-3</sup> year<sup>-1</sup> (2)



- (c) What can you conclude from the graph above about the effect of ageing on the mean concentration of protein in the blood plasma in males and females?

-----  
-----  
-----  
-----  
-----  
-----

(2)

- (d) The scientists measured the absorption of each sample of blood plasma using a colorimeter. They used a calibration curve to find the concentration of protein in samples of blood plasma.

Describe how the scientists could obtain data to produce a calibration curve and how they would use the calibration curve to find the concentration of protein in a sample of blood plasma.

-----  
-----  
-----  
-----  
-----  
-----  
-----  
-----

(3)

- (e) Older people are more likely to suffer from infectious diseases.

Suggest how this may be linked to the decrease in the mean concentration of protein in the blood as people get older.

-----  
-----  
-----

(1)

(Total 9 marks)



**Q8.**

(a) Give **three** properties of water that are important in biology. **(NOT EDEXCEL)**

1. ....
2. ....
3. ....

**(3)**

A student investigated the effect of different concentrations of sucrose solution on "chips" cut from a potato. Each chip had the same dimensions.

The student:

- weighed each chip at the start
- placed each chip in a separate test tube, each containing 10 cm<sup>3</sup> of sucrose solution at a different concentration
- left the chips in the sucrose solution for 24 hours
- dried the surface of the chips and then weighed them again.

The table shows the student's results.

Concentration of sucrose solution / mol dm <sup>-3</sup>	Initial mass of chip / g	Final mass of chip / g	Ratio of final mass to initial mass of chips
0.0	2.79	3.82	
0.2	2.75	2.97	
0.4	2.78	2.67	
0.6	2.69	2.31	
0.8	2.72	2.20	
1.0	2.77	1.99	

(b) The student produced the sucrose solutions with different concentrations from a concentrated sucrose solution.


Name the method she would have used to produce these sucrose solutions.

Name of method .....

**(1)**



- (c) Calculate the ratio of final mass to initial mass of potato chips and plot a suitable graph of your processed data. Express the ratios in the table in part (a) as a single number (for example 5.26:1 would be expressed as 5.26).



(3)

- (d) Explain the result for the chip in  $0.8 \text{ mol dm}^{-3}$  sucrose solution.

-----

-----

-----

-----

-----

-----

(2)

**(Total 9 marks)**



Q9.

(a) Describe the induced-fit model of enzyme action.

-----

-----


-----

-----

(2)

(b) A scientist investigated the hydrolysis of starch. He added amylase to a suspension of starch and measured the concentration of maltose in the reaction mixture at regular intervals.

His results are shown in the graph below.



Determine the rate of the reaction **at** 10 minutes. Show how you obtained your answer.

Rate of reaction ..... mg dm<sup>-3</sup> min<sup>-1</sup>

(2)

(c) Explain the results shown in the graph.

-----

-----

-----

-----

(2)



- (d) A quantitative Benedict's test produces a colour whose intensity depends on the concentration of reducing sugar in a solution. A colorimeter can be used to measure the intensity of this colour.

The scientist used quantitative Benedict's tests to produce a calibration curve of colorimeter reading against concentration of maltose.

Describe how the scientist would have produced the calibration curve and used it to obtain the results in the graph.

Do **not** include details of how to perform a Benedict's test in your answer.

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

-----

(3)  
(Total 9 marks)