



Topics Covered:

Introduction Why Use The Window of Tolerance? Presence in Sessions Tracking Resiliency


A Natural Processing Training Production © 2019 Craig Penner





Presence

One of the fundamental factors that determines the effectiveness of a therapy session is the client's ability to be present.




Part One - Using the Window of Tolerace, with Craig Penner


If a client is not "present" in the session, then growth and change cannot occur.





Equally important is the total presence of the therapist, which is not to be taken for granted.















Overview of this training-17 Topics

6. Therapist need to be in WoT to be able to read if the client is "in the window." *Are we both in the window?*

7. Learn to notice more, guess less. This helps therapist to be more present, and less interpretative.

8. Problem of "top down" cognitive questioning and reporting. What happens if you are NOT tracking resiliency?


9. "Faux" WoT and overriding





Overview of this training-17 Topics

14. Why SUDS reports and tracking resiliency somatically give very different data.

15. How somatic tracking tells us when the focus needs to be on helping the client back in the Window of Tolerance; dynamics of *how* clients go into hyper- and hypo-aroused states, and their abilities to regain presence.








I tell them that a lot of the reactions we all have are driven by more primitive parts of the brain, such as the limbic and reptilian brain, that do not have verbal language.





"This approach may help you with the situations, thoughts, and feelings that have been difficult. And a helpful way to look at this is to get in touch with your body as we go."



Part One - Using the Window of Tolerace, with Craig Penner




"This helps us both tune in to what is difficult and what needs working through."

You can use the Window of Tolerance as a handout with your clients.

"We can all find ourselves on here" and go over the different presentations that show up on a nervous system level.

	Window of Tolerance	
	Panic	
	Feeling overwhelmed	
	Can't think clearly	
	Emotional reactivity	
	Hyper-vigilance	
	Intrusive imagery Tension and rapid breathing	
	Increased response to sensation; startle	
	Increased least rate	
Hyper-Arousal		
		"Faux WoT"
		"Faux WoT"
Window	v of Tolerance	"Faux WoT"
	/ of Tolerance Arousal Zone	"Faux WoT"
Optima		"Faux WoT"
Optima	Arousal Zone	"Faux WoT"
Optima	Arousal Zone	"Faux WoT"
Optima Workab	Arousal Zone	
Optima	Arousal Zone le range of resiliency	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate Relative absence of awareness of sensation	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate Relative absence of awareness of sensation Numbing of emotions	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate Relative absence of awareness of sensation Numbing of emotions Disabled cognitive processing	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate Relative absence of awareness of sensation Numbing of emotions Disabled cognitive processing Reduced physical movement	
Optima Workab	Arousal Zone le range of resiliency Decreased heart rate Relative absence of awareness of sensation Numbing of emotions Disabled cognitive processing	






"Is any of this familiar for you?

There is an "o	ptimal range" in the n	niddle.
Hyper-Arousal	Increased response to sensation; startle Increased heart rate	
		"Faux WoT"
Optimal A	of Tolerance Arousal Zone a range of resiliency	
		"Faux WoT"
Hypo-Arousal	Decreased heart rate Relative absence of awareness of sensation	
	Part One - Using the Window of Tolerace, with Craig Penner	



Hypo-Arousal and the WoT


- Decrease heart rate
- Absence of sensation
- Numbing of emotions
- Movement slows or stops
- Hard to track conversation
- Poor contact with you, less contact with themselves




It's not just your cognitive understanding, it's not just your thoughts that will help get you through this ... it's also about how the rest of you works.

Your nervous system can take you to a place you didn't intend to go.







If the client cannot give specifics then they are likely not present.

Therapist can make the dynamic of "not answering" explicit.



Part One - Using the Window of Tolerace, with Craig Penner

Asking for details can help slow things down.

This helps reveal underlying processes and facilitates selfreflection.



